



University of Groningen

Efficacy of implantable cardioverter defibrillators in adults with congenital heart disease

Koyak, Zeliha; De Groot, Joris R.; Van Gelder, Isabelle C.; Budts, Werner; Van Erven, Lieselot; Van Dessel, Pascal F. H. M.; Van Dijk, Arie; Smeets, Joep L. R. M.; Maass, Alexander H.; Wilde, Arthur A. M.

Published in:
Journal of the American College of Cardiology

DOI:
[10.1016/S0735-1097\(11\)60407-5](https://doi.org/10.1016/S0735-1097(11)60407-5)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2011

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Koyak, Z., De Groot, J. R., Van Gelder, I. C., Budts, W., Van Erven, L., Van Dessel, P. F. H. M., ... Mulder, B. J. M. (2011). Efficacy of implantable cardioverter defibrillators in adults with congenital heart disease. *Journal of the American College of Cardiology*, 57(14), E407. [https://doi.org/10.1016/S0735-1097\(11\)60407-5](https://doi.org/10.1016/S0735-1097(11)60407-5)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



CONGENITAL CARDIOLOGY SOLUTIONS (ADULT CONGENITAL AND PEDIATRIC CARDIOLOGY)

EFFICACY OF IMPLANTABLE CARDIOVERTER DEFIBRILLATORS IN ADULTS WITH CONGENITAL HEART DISEASE

ACC Oral Contributions

Ernest N. Morial Convention Center, Room 225

Monday, April 04, 2011, 2:00 p.m.-2:15 p.m.

Session Title: Outcomes in Congenital Heart Disease

Abstract Category: 43. Adult Congenital Heart Disease

Presentation Number: 913-3

Authors: *Zeliha Koyak, Joris R. De Groot, Isabelle C. Van Gelder, Werner Budts, Lieselot Van Erven, Pascal F.H.M Van Dessel, Arie Van Dijk, Joep L.R.M. Smeets, Alexander H. Maass, Arthur A.M. Wilde, Petronella G. Pieper, Barbara J.M. Mulder, Academic Medical Centre, Amsterdam, The Netherlands*

Background: To date, little is known about the value of implantable cardioverter defibrillators (ICD) in adults with congenital heart disease (CHD). We investigated the long term outcomes of ICD therapy in adults with CHD.

Methods: One-hundred thirty seven adults with CHD (mean age 40, 66% male) and ICD were identified from 9 centers in the Netherlands and Belgium. ICDs were implanted for primary prevention in 70 (51%) and secondary prevention in 67 (49%) patients. Diagnosis included tetralogy of Fallot (50%), septal defects (18%), (cc) transposition of the great arteries (12%), univentricular heart (7%), and other (13%). Patients with additional primary electrical disease or cardiomyopathy were excluded.

Results: Overall, 20 patients died (9 in the primary, 11 in the secondary prevention group) during a median follow-up of 4.6 years (range 1-13 years). The cause of death was heart failure (40%), sustained ventricular tachycardia (VT) (10%), sudden death (5%), non-cardiac death (20%) and unknown causes (25%). Thirteen patients (19%) received ≥ 1 appropriate shock in the primary prevention group and 27 patients (40%) in the secondary prevention group ($P=0.004$). Overall, secondary prevention (OR 3.4 $P=0.03$), and follow up duration (OR 1.2, $P=0.005$) predicted appropriate shocks. Independent predictors of appropriate shocks were documented nonsustained VT in the primary prevention group (OR 4.7, $P=0.02$) and any VT ablation in the secondary prevention group (OR 6.8, $P=0.02$). Inappropriate shocks (mean 5.0, range 1-27) occurred in 40 patients (29%). Patients with inappropriate shocks significantly underwent more ablations for supraventricular tachycardia (OR 3.0, $P=0.01$). Additionally, 36 patients (26%) experienced 51 implantation related complications: 12 early (24%) and 39 (76%) late of which 24 (47%) lead failures.

Conclusions: Adults with CHD experience a high rate of appropriate ICD shocks related to secondary prevention and longer follow-up duration. The high efficacy rate of ICDs however, is accompanied by a high rate of procedure related complications and inappropriate shocks.